Please check the examination details below before entering your candidate information			
Candidate surname		Other names	
	Centre Number	Candidate Numbe	
Pearson Edexcel	Centre Number	Candidate Numbe	
Level 3 GCE			
Thursday 11 L	una 20	20	
Thursday 11 June 2020			
Afternoon (Time: 1 hour 30 minute	s) Paper R	reference <b>9FM0/3C</b>	
Fruther Mathematics			
Further Mathem	atics		
Further Mathem	natics		
Advanced			
Advanced		(Total Ma	

Candidates may use any calculator permitted by Pearson regulations.
Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

## **Instructions**

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
   there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Unless otherwise indicated, whenever a numerical value of g is required, take  $g = 9.8 \,\mathrm{m \, s^{-2}}$  and give your answer to either 2 significant figures or 3 significant figures.

## Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 7 questions in this question paper. The total mark for this paper is 75.
- The marks for **each** question are shown in brackets
  - use this as a guide as to how much time to spend on each guestion.

## Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶







1.	A particle $P$ of mass 0.5 kg is moving with velocity $(4\mathbf{i} + 3\mathbf{j}) \mathrm{m} \mathrm{s}^{-1}$ when it receives an impulse $\mathbf{J} \mathrm{N} \mathrm{s}$ . Immediately after receiving the impulse, $P$ is moving with velocity $(-\mathbf{i} + 6\mathbf{j}) \mathrm{m} \mathrm{s}^{-1}$ .	
	(a) Find the magnitude of <b>J</b> .	(4)
	The angle between the direction of the impulse and the direction of motion of $P$ immediately before receiving the impulse is $\alpha^{\circ}$	
	(b) Find the value of $\alpha$	(2)
		(3)

Question 1 continued	
(Tai	tal for Question 1 is 7 marks)
(10)	mi ioi Question i is / mains)



2. A truck of mass 1200 kg is moving along a straight horizontal road.

At the instant when the speed of the truck is  $v \, \text{m s}^{-1}$ , the resistance to the motion of the truck is modelled as a force of magnitude  $(900 + 9v) \, \text{N}$ .

The engine of the truck is working at a constant rate of 25 kW.

(a) Find the deceleration of the truck at the instant when v = 25

**(4)** 

Later on, the truck is moving up a straight road that is inclined at an angle  $\theta$  to the horizontal, where  $\sin \theta = \frac{1}{20}$ 

At the instant when the speed of the truck is  $v \, \text{m s}^{-1}$ , the resistance to the motion of the truck from non-gravitational forces is modelled as a force of magnitude  $(900 + 9v) \, \text{N}$ .

When the engine of the truck is working at a constant rate of 25 kW the truck is moving up the road at a constant speed of  $V \, \text{m s}^{-1}$ .

(b) Find the value of V.

1	5	1
l	J	,
•		,

(3)
<u></u>

Question 2 continued



Question 2 continued		

Question 2 continued	
(Total for Questi	on 2 is 9 marks)
, , ,	,



3.	Two particles, $A$ and $B$ , have masses $3m$ and $4m$ respectively. The particles are moving in same direction along the same straight line on a smooth horizontal surface when they coll directly. Immediately before the collision the speed of $A$ is $2u$ and the speed of $B$ is $u$ .	
	The coefficient of restitution between $A$ and $B$ is $e$ .	
	(a) Show that the direction of motion of each of the particles is unchanged by the	
	collision.	(8)
	After the collision with $A$ , particle $B$ collides directly with a third particle, $C$ , of mass $2m$ , which is at rest on the surface.	,
	The coefficient of restitution between $B$ and $C$ is also $e$ .	
	(b) Show that there will be a second collision between A and B.	
		(6)

Question 3 continued



Question 3 continued		

Question 3 continued	
	(Total for Question 3 is 14 marks)
	(20mi ior Vaccion o 15 1 i mai k5)



**4.** [*In this question,* **i** *and* **j** *are perpendicular unit vectors in a horizontal plane.*]




Figure 1

Figure 1 represents the plan view of part of a smooth horizontal floor, where AB represents a fixed smooth vertical wall.

A small ball of mass 0.5 kg is moving on the floor when it strikes the wall.

Immediately before the impact the velocity of the ball is  $(7\mathbf{i} + 2\mathbf{j}) \,\mathrm{m}\,\mathrm{s}^{-1}$ .

Immediately after the impact the velocity of the ball is (i + 6j) m s<sup>-1</sup>.

The coefficient of restitution between the ball and the wall is *e*.

(a) Show that AB is parallel to  $(2\mathbf{i} + 3\mathbf{j})$ .

(4)

(b) Find the value of *e*.

**(5)** 

Question 4 continued	



Question 4 continued

Question 4 continued	
	Total for Question 4 is 9 marks)



5.	A smooth uniform sphere $P$ has mass 0.3 kg. Another smooth uniform sphere $Q$ , with the
	same radius as $P$ , has mass $0.2 \mathrm{kg}$ .

The spheres are moving on a smooth horizontal surface when they collide obliquely. Immediately before the collision the velocity of P is  $(4\mathbf{i} + 2\mathbf{j}) \,\mathrm{m} \,\mathrm{s}^{-1}$  and the velocity of Q is  $(-3\mathbf{i} + \mathbf{j}) \,\mathrm{m} \,\mathrm{s}^{-1}$ .

At the instant of collision, the line joining the centres of the spheres is parallel to i.

The kinetic energy of Q immediately after the collision is half the kinetic energy of Q immediately before the collision.

- (a) Find
  - (i) the velocity of P immediately after the collision,
  - (ii) the velocity of Q immediately after the collision,
  - (iii) the coefficient of restitution between P and Q, carefully justifying your answers.

(11)

(b) Find the size of the angle through which the direction of motion of *P* is deflected by the collision.

**(3)** 



Question 5 continued



Question 5 continued

Question 5 continued	
	(Total for Overting 5 to 14
	(Total for Question 5 is 14 marks)



**6.** A light elastic string with natural length l and modulus of elasticity kmg has one end attached to a fixed point A on a rough inclined plane. The other end of the string is attached to a package of mass m.

The plane is inclined at an angle  $\theta$  to the horizontal, where  $\tan \theta = \frac{5}{12}$ 

The package is initially held at A. The package is then projected with speed  $\sqrt{6gl}$  up a line of greatest slope of the plane and first comes to rest at the point B, where AB = 3l.

The coefficient of friction between the package and the plane is  $\frac{1}{4}$ 

By modelling the package as a particle,

(a) show that  $k = \frac{15}{26}$ 

**(6)** 

(b) find the acceleration of the package at the instant it starts to move back down the plane from the point B.

**(5)** 

Question 6 continued



Question 6 continued

Question 6 continued	
	Total for Question 6 is 11 marks)



Figure 2

Figure 2 represents the plan view of part of a horizontal floor, where AB and CD represent fixed vertical walls, with AB parallel to CD.

A small ball is projected along the floor towards wall AB. Immediately before hitting wall AB, the ball is moving with speed  $v \, \text{m s}^{-1}$  at an angle  $\alpha$  to AB, where  $0 < \alpha < \frac{\pi}{2}$ 

The ball hits wall AB and then hits wall CD.

After the impact with wall CD, the ball is moving at angle  $\frac{1}{2}\alpha$  to CD.

The coefficient of restitution between the ball and wall AB is  $\frac{2}{3}$ 

The coefficient of restitution between the ball and wall *CD* is also  $\frac{2}{3}$ 

The floor and the walls are modelled as being smooth. The ball is modelled as a particle.

(a) Show that  $\tan\left(\frac{1}{2}\alpha\right) = \frac{1}{3}$ 

**(7)** 

(b) Find the percentage of the initial kinetic energy of the ball that is lost as a result of the two impacts.

**(4)** 

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 7 continued		



Question 7 continued		

Question 7 continued		



Question 7 continued	
	(Total for Question 7 is 11 marks)
	TOTAL FOR PAPER IS 75 MARKS

